arXiv:math/0204220v1 [math.FA] 17 Apr 2002

2000]Primary: 43A15; Secondary: 20F65, 20F18

GROUP COHOMOLOGY AND L^p-COHOMOLOGY OF FINITELY GENERATED GROUPS

MICHAEL J. PULS

Abstract. Let G be a finitely generated, infinite group, let $p > 1$, and let $L^p(G)$ denote the Banach space $\{\sum_{x\in G} a_x x \mid \sum_{x\in G} |a_x|^p < \infty\}$. In this paper we will study the first cohomology group of G with coefficients in $L^p(G)$, and the first reduced L^p-cohomology space of G. Most of our results will be for a class of groups that contains all finitely generated, infinite nilpotent groups.

1. Introduction

In this paper G will always be a finitely generated, infinite group and S will always be a symmetric generating set for G. Let M be a right G-module. A 1-cocycle with values in M is a map $\delta : G \to M$ such that $\delta(gh) = (\delta(h))\, g + \delta(g)$ for any $g, h \in G$; a 1-coboundary is a 1-cocycle of the form $\delta(g) = xg - x$ for some $x \in M$ and for all $g \in G$. We denote by $Z^1(G, M)$ the vector space of all 1-cocycles and the vector space of all 1-coboundaries will be denoted by $B^1(G, M)$. The factor group $H^1(G, M) = Z^1(G, M)/B^1(G, M)$ is called the first cohomology group of G with coefficients in M. Suppose now that M is a topological vector space and that the action of G on M is continuous. Then we give $Z^1(G, M)$ the compact open topology. Assuming that M is Hausdorff, this means that $\delta_n \to \delta$ in $Z^1(G, M)$ if and only if $\delta_n(g) \to \delta(g)$ in M for all $g \in G$. In general $B^1(G, M)$ is not closed in $Z^1(G, M)$. The quotient space $\overline{H}^1(G) = Z^1(G, M)/\overline{B^1(G, M)}$, where $\overline{B^1(G, M)}$ is the closure of $B^1(G, M)$ in $Z^1(G, M)$, is called the first reduced cohomology space.

Let $\mathcal{F}(G)$ be the set of complex-valued functions on G. We may represent each f in $\mathcal{F}(G)$ as a formal sum $\sum_{x\in G} a_x x$ where $a_x \in \mathbb{C}$ and $f(x) = a_x$. For a real number $p \geq 1, L^p(G)$ will consist of those formal sums for which $\sum_{x\in G} |a_x|^p < \infty$. Let $\alpha \in \mathcal{F}(G)$ and $g \in G$, the right translation of α by g is the function defined by $\alpha_g(x) = \alpha(xg^{-1})$. Observe that if α is represented formally by $\sum_{x\in G} a_x x$, then α_g is represented by $\sum_{x\in G} a_{xg^{-1}} x$.

In this paper we will study the cohomology theories defined above for the case $M = L^p(G)$, and with G acting on $L^p(G)$ by right translations.

I would like to thank Peter Linnell for making many useful comments on an earlier version of this paper. I would also like to thank the Mathematics department at Rutgers University for the use of their library.

Date: March 22, 2002.

1991 *Mathematics Subject Classification.* [.

Key words and phrases. group cohomology, L^p-cohomology, central element of infinite order, harmonic function, continuous linear functional.

2. Preliminaries

Let $\mathbb{C}G$ be the group ring of G over $\mathbb{C}$. For $\alpha = \sum_{x\in G} a_x x \in \mathbb{C}G$ and $\beta = \sum_{x\in G} b_x x \in \mathcal{F}(G)$ we define a multiplication $\mathcal{F}(G) \times \mathbb{C}G \to \mathcal{F}(G)$ by

$$\beta * \alpha = \sum_{x,y} b_x a_y xy = \sum_{x\in G}\left(\sum_{y\in G} b_{xy^{-1}} a_y\right) x.$$

For $1 \le p \in \mathbb{R}$, let $D^p(G) = \{\beta \in \mathcal{F}(G) \mid \beta * (g-1) \in L^p(G) \text{ for all } g \in S\}$. Recall that S is a symmetric set of generators for G. We define a norm on $L^p(G)$ by $\| \alpha \|_p = \left(\sum_{x\in G} |a_x|^p\right)^{\frac{1}{p}}$, where $\alpha = \sum_{x\in G} a_x x \in L^p(G)$. Let $\beta = \sum_{x\in G} b_x x \in D^p(G)$ and let e be the identity element of G. We define a norm on $D^p(G)$ by $\| \beta \|_{D^p(G)} = \left(\sum_{g\in S} \| \beta * (g-1) \|_p^p + |\beta(e)|^p\right)^{\frac{1}{p}}$. Under this norm $D^p(G)$ is a Banach space. Let α_1 and α_2 be elements of $D^p(G)$. We will write $\alpha_1 \simeq \alpha_2$ if $\alpha_1 - \alpha_2$ is a constant function. Clearly $\simeq$ is an equivalence relation on $D^p(G)$. Identify the constant functions on G with $\mathbb{C}$. Now $D^p(G)/\mathbb{C}$ is a Banach space under the norm induced from $D^p(G)$. That is, if $[\alpha]$ is an equivalence class from $D^p(G)/\mathbb{C}$ then $\| [\alpha] \|_{D^p(G)/\mathbb{C}} = \left(\sum_{g\in S} \| \alpha * (g-1) \|_p^p\right)^{\frac{1}{p}}$. We shall write $\| \alpha \|_{D(p)}$ for $\| [\alpha] \|_{D^p(G)/\mathbb{C}}$.

Define a linear map T from $D^p(G)$ to $Z^1(G, L^p(G))$ by $(T\alpha)(g) = \alpha * (g-1)$. It was shown in [1, lemma 4.2] that $H^1(G, \mathcal{F}(G)) = 0$, so for each 1-cocycle $\delta \in Z^1(G, \mathcal{F}(G))$ there exists an $\alpha \in \mathcal{F}(G)$ such that $\delta(g) = \alpha * (g-1)$. This implies that T is onto. The kernel of T is $\mathbb{C}$, the constant functions on G. Thus $D^p(G)/\mathbb{C}$ is isometric with $Z^1(G, L^p(G))$. Now $B^1\big(G, L^p(G)\big) = T\big(L^p(G)\big)$, so we obtain the following:

(a) The first cohomology group of G with coefficients in $L^p(G)$, $H^1(G, L^p(G))$, is isomorphic with $D^p(G)/(L^p(G) \bigoplus \mathbb{C})$.

(b) The first reduced L^p-cohomology space of G, denoted by $\overline{H}^1_{(p)}(G)$, is isometric with $D^p(G)/\overline{L^p(G) \bigoplus \mathbb{C}}$, where the closure is taken in $D^p(G)$.

3. A sufficient condition for the vanishing of $\overline{H}^1_{(p)}(G)$

In this section we give sufficient conditions on $L^p(G)$ so that $\overline{H}^1_{(p)}(G) = 0$. We begin with

Lemma 3.1. *Let $1 \le p < \infty$ and let $\alpha \in D^p(G)/\mathbb{C}$ be a non-negative, real-valued function. If $\{\beta_n\}$ is a sequence in $D^p(G)/\mathbb{C}$ such that $\beta_n \ge 0$ on G, $\{\beta_n\}$ converges pointwise to ∞ and $\| \beta_n \|_{D(p)} \to 0$ as $n \to \infty$, then $\| \alpha - \min(\alpha, \beta_n) \|_{D(p)} \to 0$ as $n \to \infty$.*

Proof. Let $\alpha_n = \min(\alpha, \beta_n), U_n = \{x \in G \mid \alpha(x) > \beta_n(x)\}$ and $V_n = \{x \mid x \in U_n \text{ or } xg^{-1} \in U_n \text{ for some } g \in S\}$. Represent α by $\sum_{x\in G} a_x x, \alpha_n$ by $\sum_{x\in G} (\tilde{a}_x)_n x$

and β_n by $\sum_{x\in G}(b_x)_n x$. Now

$$\begin{aligned}\| \alpha - \alpha_n \|^p_{D(p)} &= \sum_{g\in S} \| (\alpha - \alpha_n) * (g-1) \|^p_p \\ &= \sum_{g\in S}\sum_{x\in G} \big|(a_{xg^{-1}} - a_x) - ((\tilde{a}_{xg^{-1}})_n - (\tilde{a}_x)_n)\big|^p \\ &\le \sum_{g\in S}\sum_{x\in V_n} \big(|a_{xg^{-1}} - a_x| + |(b_{xg^{-1}})_n - (b_x)_n|\big)^p \\ &\le 2^p \sum_{g\in S}\sum_{x\in V_n} \big(|a_{xg^{-1}} - a_x|^p + |(b_{xg^{-1}})_n - (b_x)_n|^p\big) \\ &= 2^p \sum_{g\in G}\big(\sum_{x\in V_n} |a_{xg^{-1}} - a_x|^p + \sum_{x\in V_n} |(b_{xg^{-1}})_n - (b_x)_n|^p\big).\end{aligned}$$

Let $\epsilon > 0$ be given. Since $\alpha \in D^p(G)/\mathbb{C}$ there exists a finite subset F of G such that $\sum_{g\in S}\sum_{x\in G\setminus F} |a_{xg^{-1}} - a_x|^p < \epsilon$. Since $\beta_n(x) \to \infty$ as $n \to \infty$, there exists N such that $V_n \subseteq G \setminus F$ for all $n \ge N$. Thus $\sum_{g\in S}\sum_{x\in V_n} |a_{xg^{-1}} - a_x|^p \to 0$ as $n \to \infty$. By hypothesis $\sum_{g\in S}\sum_{x\in V_n} |(b_{xg^{-1}})_n - (b_x)_n|^p \to 0$ as $n\to\infty$. Therefore $\| \alpha - \min(\alpha, \beta_n) \|_{D(p)} \to 0$ as $n \to \infty$. □

Let $\alpha \in D^p(G)$ and $x \in G$. Then $\big|\alpha(x)\big|$ will denote the modulus of $\alpha(x)$ and $|\alpha|$ will denote the function $\big|\alpha\big|(x) = \big|\alpha(x)\big|$. We are now ready to give a sufficient condition for the vanishing of $\overline{H}^1_{(p)}(G) = 0$.

Theorem 3.2. *Let $1 \le p < \infty$. Suppose there exists a sequence $\{\alpha_n\}$ in $L^p(G)$ such that $\| \alpha_n \|_{D(p)} \to 0$ as $n \to \infty$ and $\{\alpha_n(x)\}$ does not converge pointwise to zero for each x in G. Then $\overline{H}^1_{(p)}(G) = 0$.*

Proof. By taking a subsequence if necessary we may assume that $\| \alpha_n \|_{D(p)} < \frac{1}{n^2}$ for all n. Since $\| |\alpha| \|_{D(p)} \le \| \alpha \|_{D(p)}$ we may assume that $\alpha_n(x) \ge 0$ for all $x \in G$. Set $\beta_n = n\alpha_n$. Now $\beta_n(x) \ge 0$ for all $x \in G, \beta_n(x) \to \infty$ as $n \to \infty$ for every $x \in G$ and $\| \beta_n \|^p_{D(p)} = \| n\alpha_n \|^p_{D(p)} = n^p \| \alpha_n \|^p_{D(p)} \le n^p \left(\frac{1}{n^{2p}}\right) = \frac{1}{n^p}$. We now have that $\| \beta_n \|_{D(p)} \to 0$ as $n \to \infty$. Let α be a real-valued, non-negative function in $D^p(G)/\mathbb{C}$. By Lemma 3.1, $\| \alpha - \min(\alpha, \beta_n) \|_{D(p)} \to 0$ as $n \to \infty$. Thus $\alpha \in \overline{L^p(G)}$ since $\min(\alpha, \beta_n) \in L^p(G)$. It now follows by approximation that $\overline{L^p(G)} = D^p(G)/\mathbb{C}$. Hence $\overline{H}^1_{(p)}(G) = D^p(G)/\overline{(L^p(G) \bigoplus \mathbb{C})} = 0$. □

4. Harmonic Functions

In this section we will give some results about harmonic functions on G. Let $\alpha \in \mathcal{F}(G)$ and represent α by $\sum_{x\in G} a_x x$. Now define

$$(\triangle\alpha)(x) := \sum_{g\in S}\big((\alpha * (g-1))(x)\big) = \sum_{g\in S}(a_{xg^{-1}} - a_x).$$

We shall say that α is harmonic on G if $(\triangle\alpha)(x) = 0$ for each $x \in G$, alternatively α is harmonic if $|S|\big(\alpha(x)\big) = \sum_{g\in S} \alpha(xg^{-1})$ for each x in G. Let $LHD^p(G) = \{\alpha \mid \alpha$ is harmonic and $\alpha \in D^p(G)\}$. Observe that the constant functions are contained in $LHD^p(G)$.

Lemma 4.1. *Let $x \in G$. There exists a positive constant M_x such that $|\alpha(x)| \le M_x \| \alpha \|_{D^p(G)}$ for all $\alpha \in D^p(G)$.*

Proof. Write $x = g_1 g_2 \dots g_n$ where $g_k \in S$ and no subword of $g_1 g_2 \dots g_n$ is the identity. Set $w_k = g_1 g_2 \dots g_k$. Let $\alpha \in D^p(G)$. Now

$$\begin{aligned}|\alpha(x)| &= \big(|\alpha(w_n) - \alpha(w_{n-1}) + \alpha(w_{n-1}) - \alpha(w_{n-2}) + \\ &\quad \cdots + \alpha(w_2) - \alpha(w_1) + \alpha(w_1) - \alpha(e) + \alpha(e)|^p\big)^{\frac{1}{p}} \\ &\le \big((|\alpha(w_n) - \alpha(w_{n-1})| + |\alpha(w_{n-1}) - \alpha(w_{n-2})| + \\ &\quad \cdots + |\alpha(w_2) - \alpha(w_1)| + |\alpha(w_1) - \alpha(e)| + |\alpha(e)|)^p\big)^{\frac{1}{p}}.\end{aligned}$$

If $0 \le a_1, \dots, a_n \in \mathbb{R}$, then by Jensen's inequality [3, p. 189] applied to the function x^p for $x > 0$,

$$(a_1 + \cdots + a_n)^p \le n^{p-1}(a_1^p + \cdots + a_n^p),$$

consequently

$$\begin{aligned}|\alpha(x)| &\le \big(n^{p-1}(|\alpha(w_n) - \alpha(w_{n-1})|^p + |\alpha(w_{n-1}) - \alpha(w_{n-2})|^p + \\ &\quad \cdots + |\alpha(w_1) - \alpha(e)|^p + |\alpha(e)|^p)\big)^{\frac{1}{p}} \\ &= n^{\frac{p-1}{p}}\big(|(\alpha * (g_n^{-1} - 1))(w_{n-1})|^p + |(\alpha * (g_{n-1}^{-1} - 1))(w_{n-2})|^p + \\ &\quad \cdots + |(\alpha * (g_1^{-1} - 1))(e)|^p + |\alpha(e)|^p\big)^{\frac{1}{p}} \\ &\le n^{\frac{p-1}{p}} \parallel \alpha \parallel_{D^p(G)} .\end{aligned}$$

□

We are now ready to prove:

Lemma 4.2. *The set $LHD^p(G)$ is closed in $D^p(G)$.*

Proof. Let $\{\alpha_n\}$ be a sequence in $LHD^p(G)$ and suppose that $\{\alpha_n\} \to \alpha$ in $D^p(G)$. Let $x \in G$. By Lemma 4.1 there exists a positive constant M_x such that $|(\alpha - \alpha_n)(x)| \le M_x \parallel \alpha - \alpha_n \parallel_{D^p(G)}$. Thus $\{\alpha_n(x)\}$ converges pointwise to $\alpha(x)$ for all $x \in G$. Represent α by $\sum_{x \in G} a_x x$ and α_n by $\sum_{x \in G} (\tilde{a}_x)_n x$. Now $\sum_{g \in S}\big((\tilde{a}_{xg^{-1}})_n - (\tilde{a}_x)_n\big) = 0$ for all natural numbers n and for all $x \in G$. Thus $\sum_{g \in S}(a_{xg^{-1}} - a_x) = 0$ for all $x \in G$. □

The following proposition will be used in sections 5 and 6. The idea behind the proposition and the proof were inspired by [4, Theorem 3.1].

Proposition 4.3. *Let G be a finitely generated group that has a central element of infinite order. If $1 \le p < \infty$, then $LHD^p(G) = \mathbb{C}$.*

Proof. Let y be an element of infinite order that is an element of the center of G. Let $\alpha \in LHD^p(G)$ and $x \in G$. Now $|S|\big(\alpha_y(x)\big) = |S|\big(\alpha(xy^{-1})\big) = \sum_{g \in S} \alpha(xy^{-1}g^{-1}) = \sum_{g \in S} \alpha(xg^{-1}y^{-1}) = \sum_{g \in S} \alpha_y(xg^{-1})$. Hence α_y is harmonic if α is harmonic. Define a new function $\beta(x) = \alpha_y(x) - \alpha(x)$ on G. Now β is harmonic since it is the sum of harmonic functions. The formal series representation of β is $\sum_{x \in G}(a_{xy^{-1}} - a_x)x$. Since $\alpha \in D^p(G)$ we have that $\beta \in L^p(G)$. Thus for each $\epsilon > 0$, the set $\{x \mid |a_{xy^{-1}} - a_x| > \epsilon\}$ is finite. By the maximum (minimum) principle for harmonic functions it must be the case $|a_{xy^{-1}} - a_x| < \epsilon$ for all $x \in G$. Hence $\alpha_y(x) = a_{xy^{-1}} = a_x = \alpha(x)$ for all $x \in G$.

Let $x \in G$ and $g \in S$. We now have that $\alpha(x) - \alpha(xg) = \alpha_y(x) - \alpha_y(xg) = \alpha_{y^2}(x) - \alpha_{y^2}(xg) = \cdots = \alpha_{y^n}(x) - \alpha_{y^n}(xg)$ for all natural numbers n. In other words , $a_x - a_{xg} = a_{xy^{-1}} - a_{xy^{-1}g} = \cdots = a_{xy^{-n}} - a_{xy^{-n}g}$. Since $\alpha \in D^p(G)$ and $y^n \neq y$ for all natural numbers n, we have that $|a_{xy^{-n}} - a_{xy^{-n}g}| < \epsilon$ for all $\epsilon > 0$. Thus $\alpha(x) = \alpha(xg)$. The proposition now follows since S generates G. ☐

Remark 4.4. The center of a finitely generated, infinite nilpotent group contains an element of infinite order.

5. Groups with a central element of infinite order.

Let $1 < p \in \mathbb{R}$ and let d be a natural number. It was proven in [2] that $\mathbb{Z}^d$ satisfies the hypothesis of Theorem 3.2 if and only if $d \leq p$. Thus, for example, Theorem 3.2 cannot be used to determine if $\overline{H}^1_{(p)}(\mathbb{Z}^d) = 0$ whenever $d > p$. In this section we will prove that $\overline{H}^1_{(p)}(G) = 0$ whenever G is a group that has a central element of infinite order.

Given $1 < p \in \mathbb{R}$, we shall always let q denote the conjugate index of p. Thus if $p > 1$, then $\frac{1}{p} + \frac{1}{q} = 1$. Fix $\beta = \sum_{x \in G} b_x x \in D^q(G)/\mathbb{C}$. We can define a linear functional on $D^p(G)/\mathbb{C}$ by $\langle \alpha, \beta \rangle = \sum_{x \in G} \sum_{g \in S} \big((\alpha * (g-1))(x)\big)\big(\overline{(\beta * (g-1))(x)}\big) = \sum_{x \in G} \sum_{g \in S} (a_{xg^{-1}} - a_x)(\overline{b_{xg^{-1}} - b_x})$, where $\alpha = \sum_{x \in G} a_x x \in D^p(G)/\mathbb{C}$. The sum is finite since $\alpha * (g-1) \in L^p(G)$ and $\beta * (g-1) \in L^q(G)$ for each $g \in S$. For $y \in G$, define δ_y by $\delta_y(x) = 0$ if $x \neq y$ and $\delta_y(y) = 1$.

Lemma 5.1. *Let $\alpha \in \mathcal{F}(G)$. Then α is a harmonic function if and only if $\langle \delta_y, \alpha \rangle = 0$ for all $y \in G$.*

Proof. Represent α by $\sum_{x \in G} a_x x$ and let $y \in G$. Now

$$\langle \delta_y, \alpha \rangle = -2 \sum_{g \in S} (\overline{a_{yg^{-1}} - a_y}).$$

If α is harmonic, then $\langle \delta_y, \alpha \rangle = 0$. Conversely, if $\langle \delta_y, \alpha \rangle = 0$ for all $y \in G$, then α is harmonic since $\sum_{g \in S} (a_{yg^{-1}} - a_y) = 0$ for all $y \in G$. ☐

For $X \subseteq D^p(G)/\mathbb{C}$, let $\big(\overline{X}\big)_{D(p)}$ denote the closure of X in $D^p(G)/\mathbb{C}$.

Proposition 5.2. *If $\alpha \in \big(\overline{\mathbb{C}G}\big)_{D(p)}$ and $\beta \in LHD^q(G)$, then $\langle \alpha, \beta \rangle = 0$.*

Proof. Let $\{\alpha_n\}$ be a sequence in $\mathbb{C}G$ which converges to α in $D^p(G)/\mathbb{C}$. It follows from Lemma 5.1 that $\langle \alpha_n, \beta \rangle = 0$ for each n. We now obtain,

$$\begin{aligned}
0 &\leq \left| \sum_{x \in G} \sum_{g \in S} \big((\alpha * (g-1))(x)\big)\big(\overline{(\beta * (g-1))(x)}\big) \right| \\
&= \left| \sum_{x \in G} \sum_{g \in S} \big(((\alpha - \alpha_n) * (g-1))(x)\big)\big(\overline{(\beta * (g-1))(x)}\big) \right| \\
&\leq \sum_{x \in G} \sum_{g \in S} \big| \big(((\alpha - \alpha_n) * (g-1))(x)\big)\big(\overline{(\beta * (g-1))(x)}\big) \big| \\
&\leq \| \alpha - \alpha_n \|_{D(p)} \| \beta \|_{D(q)} \to 0 \text{ as } n \to \infty.
\end{aligned}$$

The last inequality follows from Hölder's inequality. ☐

We are now ready to prove

Theorem 5.3. *If $1 < p \in \mathbb{R}$ and G is a finitely generated group with a central element of infinite order, then $\overline{H}^1_{(p)}(G) = 0$.*

Proof. The space of continuous linear functionals on $D^p(G)/\mathbb{C}$ is $D^q(G)/\mathbb{C}$. Let $\left(\overline{L^p(G)}\right)^{\perp}_{D(p)} = \{\beta \in D^q(G)/\mathbb{C} \mid \langle \alpha, \beta \rangle = 0 \text{ for all } \alpha \in \left(\overline{L^p(G)}\right)_{D(p)}\}$. Since $\left(\overline{\mathbb{C}G}\right)_{D(p)} = \left(\overline{L^p(G)}\right)_{D(p)}$ it follows from Proposition 5.2 that $LHD^q(G)/\mathbb{C}$ is contained in $\left(\overline{L^p(G)}\right)^{\perp}_{D(p)}$.

Let $\beta \in D^q(G)/\mathbb{C}$ and represent β by $\sum_{x \in G} b_x x$. Suppose that β is not harmonic on G. Then there exists an $x \in G$ such that $\sum_{g \in S}(b_{xg^{-1}} - b_x) \neq 0$. If α is supported only on x, then by Lemma 5.1 we have that $\langle \alpha, \beta \rangle \neq 0$. Thus the space of continuous linear functionals on $D^p(G)/(\overline{L^p(G) \bigoplus \mathbb{C}})$ is $LHD^q(G)/\mathbb{C}$. The theorem now follows from Proposition 4.3. $\square$

Remark 5.4. If G is a group for which $L^2(G)$ does not satisfy the hypothesis of Theorem 3.2, then using the proof of the above theorem we can obtain the well known result $D^2(G) = (\overline{\mathbb{C}G})_{D^2(G)} \bigoplus LHD^2(G)$.

6. A description of $H^1(G, L^2(G))$

Let $d > 1$. We shall say that G satisfies condition S_d if there exists a constant $C > 0$ such that $\parallel \alpha \parallel_{\frac{d}{d-1}} \leq C \parallel \alpha \parallel_{D(1)}$ for all $\alpha \in \mathbb{C}G$. In this section we will describe the nonzero elements of $H^1\big(G, L^2(G)\big)$ for groups that satisfy property S_d and have a central element of infinite order. If $\alpha \in \mathcal{F}(G)$ and $t \geq 1$, then α^t will denote the function $\alpha^t(x) = \big(\alpha(x)\big)^t$. Let us start with

Lemma 6.1. *Let G be a finitely generated group and let t be a real number greater than or equal to 2. If α is a non-negative, real function in $\mathcal{F}(G)$, then*

$$\parallel \alpha^t \parallel_{D(1)} \leq 2t \sum_{x \in G} \alpha^{t-1}(x) \left(\sum_{g \in S} \left| \big(\alpha * (g-1)\big)(x) \right| \right).$$

Proof. Represent α by $\sum_{x \in G} a_x x$. Let $x \in G$ and let $g \in S$. It follows from the Mean Value Theorem applied to x^t that $(r^t - s^t) \leq t(r^{t-1} + s^{t-1})(r - s)$ where r and s are real numbers with $0 \leq s \leq r$. Thus $|a^t_{xg^{-1}} - a^t_x| \leq t(a^{t-1}_x + a^{t-1}_{xg^{-1}})|a_{xg^{-1}} - a_x|$. Now $\parallel \alpha^t \parallel_{D(1)} = \sum_{x \in G} \sum_{g \in S} \left|\big(\alpha^t * (g-1)\big)(x)\right| = \sum_{x \in G} \sum_{g \in S} |a^t_{xg^{-1}} - a^t_x| \leq t \sum_{x \in G} \sum_{g \in S} (a^{t-1}_x + a^{t-1}_{xg^{-1}})|a_{xg^{-1}} - a_x| = 2t \sum_{x \in G} \sum_{g \in S} a^{t-1}_x |a_{xg^{-1}} - a_x|$. $\square$

We will now use this lemma to prove the following:

Proposition 6.2. *Let $d > 2$. If G satisfies condition S_d, then there is a constant $C' > 0$ such that $\parallel \alpha \parallel_{\frac{2d}{d-2}} \leq C' \parallel \alpha \parallel_{D(2)}$ for all $\alpha \in \mathbb{C}G$.*

Proof. Put $t = \frac{2d-2}{d-2}$ and represent α by $\sum_{x \in G} a_x x$. By property S_d, Lemma 6.1 and Schwartz's inequality we have (assuming without loss of generality that α is non-negative).

$$\begin{aligned}
\| \alpha^{\frac{2d-2}{d-2}} \|_{\frac{d}{d-1}} &\leq C \| \alpha^{\frac{2d-2}{d-2}} \|_{D(1)} \\
&\leq 2C(\frac{2d-2}{d-2}) \sum_{x \in G} \alpha^{\frac{d}{d-2}}(x)(\sum_{g \in S} |\big(\alpha * (g-1)\big)(x)|) \\
&= 2C(\frac{2d-2}{d-2}) \sum_{x \in G} \sum_{g \in S} a_x^{\frac{d}{d-2}} |a_{xg^{-1}} - a_x| \\
&\leq 2C(\frac{2d-2}{d-2}) \| \alpha^{\frac{d}{d-2}} \|_2 \| \alpha \|_{D(2)} .
\end{aligned}$$

Observe $\| \alpha^{\frac{2d-2}{d-2}} \|_{\frac{d}{d-1}} = \| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{d-1}{d}}$ and $\| \alpha^{\frac{d}{d-2}} \|_2 = \| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{1}{2}}$. Substituting we obtain $\| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{d-1}{d}} \leq C' \| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{1}{2}} \| \alpha \|_{D(2)}$. The proposition follows by dividing both sides by $\| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{1}{2}}$ and observing that $\| \alpha^{\frac{2d}{d-2}} \|_1^{\frac{d-2}{2d}} = (\| \alpha \|_{\frac{2d}{d-2}}^{\frac{2d}{d-2}})^{\frac{d-2}{2d}}$. $\square$

If G is finitely generated, then $L^p(G) \subseteq L^{p'}(G)$ for $1 \leq p \leq p'$. If $\alpha \in L^p(G)$, then α is in the zero class of $H^1\big(G, L^p(G)\big)$. Our next result will show, for the case G a group that has a central element of infinite order and satisfies property S_d, that each nonzero class in $H^1\big(G, L^2(G)\big)$ can be represented by a function in $L^{p'}(G)$ for some fixed real number $p' > 2$.

Theorem 6.3. *If G is a finitely generated group that has a central element of infinite order and satisfies condition S_d for $d > 2$, then each nonzero class in $H^1\big(G, L^2(G)\big)$ can be represented by a function from $L^{\frac{2d}{d-2}}(G)$.*

Proof. Let 1_G denote the constant function one on G. If $1_G \in (\overline{\mathbb{C}G})_{D^2(G)}$, then there exists a sequence $\{\alpha_n\}$ in $\mathbb{C}G$ such that $\| 1_G - \alpha_n \|_{D^2(G)} \to 0$ but $\| \alpha_n \|_{\frac{2d}{d-2}} \not\to 0$ contradicting Proposition 6.2. Hence $(\overline{\mathbb{C}G})_{D^2(G)} \neq D^2(G)$. By Remark 5.4 we have the decomposition $D^2(G) = \overline{L^2(G)} \bigoplus LHD^2(G)$. By Proposition 4.3, $LHD^2(G) = \mathbb{C}$. Thus nonzero classes in $H^1\big(G, L^2(G)\big)$ can be represented by functions in $(\overline{\mathbb{C}G})_{D^2(G)} \setminus L^2(G)$. Let $\alpha \in (\overline{\mathbb{C}G})_{D^2(G)}$, so there exists a sequence $\{\alpha_n\}$ in $\mathbb{C}G$ such that $\alpha_n \to \alpha$ in the Banach space $D^2(G)$. Thus $\{\alpha_n\}$ is a Cauchy sequence in $D^2(G)$. By Proposition 6.2 $\{\alpha_n\}$ forms a Cauchy sequence in $L^{\frac{2d}{d-2}}(G)$. Now $\| \overline{\alpha} - \alpha_n \|_{\frac{2d}{d-2}} \to 0$ for some $\overline{\alpha} \in L^{\frac{2d}{d-2}}(G)$. Since L^p-convergence implies pointwise convergence $\| (\overline{\alpha} - \alpha_n) * (g-1) \|_2 \to 0$ as $n \to \infty$ for each $g \in S$. Hence $\| (\overline{\alpha} - \alpha_n) \|_{D^2(G)} \to 0$ as $n \to \infty$. Therefore $\overline{\alpha} = \alpha$ $\square$

Let A be a finite subset of G and define

$$\partial A := \{x \in A \mid \text{ there exists } g \in S \text{ with } xg \notin A\}.$$

We shall say G satisfies condition $(IS)_d$ if $|A|^{d-1} < |\partial A|^{d-1}$ for all finite subsets A of G. Varopoulous proves the following proposition on page 224 of [5].

Proposition 6.4. *A finitely generated group G satisfies the condition $(IS)_d$ for some $d \geq 1$ if and only if it satisfies condition S_d.*

Now $\mathbb{Z}^d$ satisfies condition $(IS)_d$ but does not satisfy $(IS)_{d+\epsilon}$ for any $\epsilon > 0$. We now have the following:

Corollary 6.5. *Let* $d \geq 3$. *Each nonzero class in* $H^1\big(\mathbb{Z}^d, L^2(\mathbb{Z}^d)\big)$ *can be represented by a function from* $L^{\frac{2d}{d-2}}(\mathbb{Z}^d)$.

REFERENCES

[1] C. H. Houghton. Ends of groups and the associated first cohomology groups. *J. London Math. Soc. (2)*, 6:81–92, 1972.

[2] Fumi-Yuki Maeda. A remark on parabolic index of infinite networks. *Hiroshima Math. J.*, 7(1):147–152, 1977.

[3] A. Wayne Roberts and Dale E. Varberg. *Convex functions*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1973. Pure and Applied Mathematics, Vol. 57.

[4] Paolo M. Soardi and Wolfgang Woess. Uniqueness of currents in infinite resistive networks. *Discrete Appl. Math.*, 31(1):37–49, 1991.

[5] N. Th. Varopoulos. Isoperimetric inequalities and Markov chains. *J. Funct. Anal.*, 63(2):215–239, 1985.

MATHEMATICS, NEW JERSEY CITY UNIVERSITY, 2039 KENNEDY BLVD., JERSEY CITY, NJ 07305–1597, USA

E-mail address: `mpuls@njcu.edu`